

Postoperative Instructions for Patients Undergoing Ear Surgery

1. Getting home and postoperative supervision.

It is extremely hazardous to drive after general anesthesia or intravenous sedation. Please arrange for a responsible adult to drive you home and stay with you for at least 24 hours.

2. Activity for the first 24 hours.

The effects of anesthesia can remain in your body as long as 24 hours. After surgery, plan to rest for 24 hours. Avoid straining and stooping. Sit upright as much as possible.

3. Sleeping position.

Sleeping with the head elevated on two pillows decreases the amount of bruising, swelling and pain.

4. Diet.

After surgery you may take liquids such as soup, tea, soft drinks, or Jell-O. Gradually progress first to a soft diet and then to a regular diet over the next 24 hours.

5. Medications, herbal supplements, vitamins and alcohol.

Fill your prescriptions and take your post-operative medications as directed. If you should become nauseated or develop a rash, call the office. You may resume your regular medications unless Dr. Boyd directs otherwise. However, avoid aspirin and aspirin-containing products for two weeks before and after surgery. Remember, many herbal supplements adversely affect the clotting process and actively cause bruising and bleeding. These should be stopped two weeks before surgery and may be resumed two weeks afterwards. Avoid alcohol until post-operative medications are finished. Dr. Boyd does not recommend taking mega doses of vitamins in excess of the normal daily requirement (which is more than adequately met by a normal balanced diet). Please discontinue this form of medication for two weeks before and after surgery unless you are being treated for a specific deficiency. One multivitamin tablet per day is permissible, however.

6. Dressings.

After surgery, slight spotting of blood may be seen on the bandages. However enlarging bright red blood spots or dripping blood should be reported to the office immediately. The bandages should only be removed by Dr. Boyd or one of his staff. Do not get them wet. If the bandages are too tight or too loose, call the office. The bandages are removed at one week after which a tennis headband is applied. You will be instructed in its use. Once the bandages are removed the ears may appear swollen and bluish in color. This is normal.

7. Wound care, shampooing, showering.

After the bandages are removed, wash your hair daily with a mild shampoo. Wear a tennis headband at night for six weeks to prevent damage to the repair. Your sutures (stitches) will be removed two weeks after surgery.

8. Activity after the first 24 hours (including driving).

Avoid strenuous activity for two weeks after surgery. However, a moderate amount of walking is encouraged. Do not drive for one week and never drive while taking pain medication. You may wear makeup after checking with Dr. Boyd.

9. Things to avoid. (See 2 and 5 also)

It is vital that you do not smoke after surgery to avoid respiratory and wound healing complications. Avoid touching, hitting or brushing the ears.

10. Bruising, swelling and other normal events.

A moderate amount of swelling and bruising is common. It will take several weeks before you can see the final result. If you have unusual or severe pain or swelling (especially on one side), bleeding from the dressing or any other concerns, do not hesitate to get in touch with Dr. Boyd. Someone in the office will call you the day after surgery to check on your condition and answer any questions you might have. You can reach Dr. Boyd or a staff member around the clock by calling (310) 530-4200.

These instructions are not intended to cover every possible problem that could arise. Good judgment on your part is essential for a good result.