

Postoperative Instructions for Patients Undergoing Neck Liposuction and/or Chin Augmentation

1. Getting home and postoperative supervision.

It is extremely hazardous to drive after general anesthesia or intravenous sedation. Please arrange for a responsible adult to drive you home and stay with you for at least 24 hours.

2. Activity for the first 24 hours.

The effects of anesthesia can remain in your body as long as 24 hours. After surgery, plan to rest for 24 hours. Avoid straining and stooping.

3. Sleeping position.

For three days after surgery sleep on your back elevated with 2 pillows.

4. Diet.

After surgery you may take liquids such as soup, tea, soft drinks, or Jell-O. Gradually progress to a soft diet and then to a regular diet over the next 24 hours as tolerated.

5. Medications, herbal supplements, vitamins and alcohol.

Fill your prescriptions and take your post-operative medications as directed. If you should become nauseated or develop a rash, call the office. You may resume your regular medications unless Dr. Boyd directs otherwise. However, avoid aspirin and aspirin-containing products for two weeks before and after surgery. Remember, many herbal supplements adversely affect the clotting process and actively cause bruising and bleeding. These should be stopped two weeks before surgery and may be resumed two weeks afterwards. Avoid alcohol until post-operative medications are finished. Dr. Boyd does not recommend taking mega doses of vitamins in excess of the normal daily requirement (which is more than adequately met by a normal balanced diet). Please discontinue this form of medication for two weeks before and after surgery unless you are being treated for a specific deficiency. One multivitamin tablet per day is permissible, however.

6. Dressings.

A chinstrap or tape must be worn for one week.

7. Wound care, shampooing, showering.

You may shower and shampoo.

8. Activity after the first 24 hours (including driving).

Avoid strenuous activity for two weeks after surgery. However, walking is encouraged. Do not drive for one week and never while taking pain medication. You may wear makeup after checking with Dr. Boyd.

9. Things to avoid. (See 2 and 5 also)

It is vital that you do not smoke after surgery to avoid respiratory and wound healing complications. Avoid sun exposure and wear sunscreen over the suture lines when it cannot be avoided.

10. Bruising, swelling and other normal events.

You may notice a change in sensation over the chin and lower lip. This is common and will return to normal within 2 to 4 months. The scar will be red for approximately six months and then begin to fade and soften.

A moderate amount of swelling and bruising is common. It will take several weeks before you can see the final result. If you have unusual or severe pain or swelling (especially on one side of your face or neck), bleeding from the dressing or any other concerns, do not hesitate to contact the office.

Someone in the office will call you the day after surgery to check on your condition and answer any questions you might have. You can reach Dr. Boyd or a staff member around the clock by calling (310) 295-2287.

These instructions are not intended to cover every possible problem that could arise. Good judgment on your part is essential for a good result.