

Postoperative Instructions for Patients Undergoing Thigh Reduction/Thigh Lift

1. Getting home and postoperative supervision.

It is extremely hazardous to drive after general anesthesia or intravenous sedation. Please arrange for a responsible adult to drive you home and stay with you for at least 24 hours.

2. Activity for the first 24 hours.

The effects of anesthesia can remain in your body as long as 24 hours. After surgery, plan to rest for 24 hours. Avoid straining and stooping. Generally, when resting, elevate the legs.

3. Diet.

After surgery you may take liquids such as soup, tea, soft drinks, or Jell-O. Gradually progress to a soft diet and then a regular diet over the next 24 hours.

4. Medications, herbal supplements, vitamins and alcohol.

Fill your prescriptions and take your post-operative medications as directed. If you should become nauseated or develop a rash, call the office. You may resume your regular medications unless Dr. Boyd directs otherwise. However, avoid aspirin and aspirin-containing products for two weeks before and after surgery. Remember, many herbal supplements adversely affect the clotting process and actively cause bruising and bleeding. These should be stopped two weeks before surgery and may be resumed two weeks afterwards. Avoid alcohol until post-operative medications are finished. Dr. Boyd does not recommend taking megadoses of vitamins in excess of the normal daily requirement (which is more than adequately met by a normal balanced diet). Please discontinue this form of medication for two weeks before and after surgery unless you are being treated for a specific deficiency. One multivitamin tablet per day is permissible, however.

5. Dressings.

The elastic pressure garment should remain on almost all the times for the first week. For the next three weeks it must be worn all day long and removed on retiring to bed. However, it may be removed for daily showering and replaced as quickly as possible. Leave the transparent adhesive dressing in place until Dr. Boyd removes it.

6. Wound care, shampooing, showering.

7. Two days after surgery, remove the elastic garment and any gauze, leaving the transparent dressing in place. Place Vaseline around the points where the drains exit the skin. Take a shower, but do not immerse yourself in the bathtub. Thereafter, shower and shampoo daily replacing the garment each time. Do this for one month.

8. Activity after the first 24 hours (including driving).

The day after surgery, get out of bed and walk around the house. Do this several times a day increasing the activity as tolerated. After two weeks, you may swim and walk longer distances, but avoid strenuous exercise for six weeks. Do not drive for one week and never while taking pain medication.

9. Things to avoid. (See 2 and 5 also)

It is vital that you do not smoke after surgery to avoid respiratory and wound healing complications.

10. Bruising, swelling and other normal events.

It is normal to have swelling and bruising of the surgical area. This will gradually improve over several weeks. However, fever, shortness of breath, chest pain or unusual pain in the operated area, especially on one side, should be called to the attention of Dr. Boyd immediately.

Someone in the office will call you the day after surgery to check on your condition and answer any questions you might have. You can reach Dr. Boyd or a staff member around the clock by calling (310) 295-2287.

These instructions are not intended to cover every possible problem that could arise. Good judgment on your part is essential for a good result.